

Erosion Prevention Through Empowerment of Human Resources to Support Food Security Around the Kambaniru Watershed, East Nusa Tenggara

Abstract. The high activity of the community that is not accompanied by the readiness of human resources around the watershed determines the status of the land use index, the erosion index, and economic and social factors. The negative externality impacts arise due to unfriendly land-use management on various types of cultivation, including agriculture, cultivation, plantations, or even livestock, contributing to the potential for erosion. This study aims to create a causal model of erosion prevention through a human resource development approach to support community food security and develop a technical program for land rehabilitation around the Kambaniru watershed, East Nusa Tenggara. The research method is presented descriptively with observation, tabulation, and data valuation techniques. The performance assessment results of the Kambaniru watershed show a medium category classification. Multipurpose plants and trees are prioritized for land rehabilitation to prevent erosion and provide food security for the poor, such as corn and tubers. Human resource capacity development in the upstream and central communities has successfully supported the erosion prevention and land rehabilitation program. The policy implications of this research are as follows: the limited capacity of community human resources in land conservation can be started by socializing the planting of short-lived trees. The recommended species are legume family types that can thrive in various types of soil and are flexible with climate change. Corn and tubers are a source of high calories and nutrients needed to improve the quality of human resources in the Kambaniru watershed area. Therefore, it is necessary to develop a food reserve land program in the Kambaniru watershed area.

Keywords: Human Resources Capacity, Food Security, Environmental Degradation, Watershed Health

1. Introduction

Forest village communities fulfill their daily needs by carrying out various activities in the Kambaniru Watershed (DAS). This watershed has an area of 140,440.70 hectares and is located in East Sumba Regency, East Nusa Tenggara. The Kambaniru watershed includes nine sub-districts, consisting of 34 villages and eight sub-districts. The Kambaniru dam uses water from this watershed to irrigate 1,540 hectares of rice fields [1].

Environmental degradation causes forest cover in watershed areas to decrease drastically. This condition is caused by the limited capacity of community human resources in managing and utilizing land and forest resources. Freelancing farming systems, shifting ground, burning savanna during the dry season, and taking wood that is not balanced with replanting accelerate the rate of environmental degradation. This condition is exacerbated by limited access to capital support. The community, especially the poor, women, and vulnerable groups, ends up relying on forest products for a living because they do not have other business alternatives. During the last ten years, conditions in the Kambaniru watershed area have been quite alarming. In the dry season, the water discharge of the Kambaniru river decreases, but in the rainy season, it often overflows to inundate residential areas in certain places. Njurumana [2] stated that the Kambaniru watershed is experiencing severe environmental degradation but continues to be rehabilitated. Inland rehabilitation activities in the Kambaniru watershed area, there are short-lived plants. Calliandra (*Calliandra calothyrsus*) and Sengon (*Albizia falcataria*) plants are used for reforestation and preventing erosion. According to Krisnawati *et al.* [3]; Hashim *et al.* [4]; and Angima *et al.* [5], both are tree plants from the leguminous family type that can thrive in various types of soil and are flexible with climate and climate change.

The high activity of the community around the watershed sometimes also determines the poor land use index (IPL), erosion index, and economic and social factors. The negative external impacts arising from land-use activities that are not environmentally friendly in every type of cultivation, whether agriculture, cultivation, plantations or even livestock, contribute to the potential for erosion. The pattern of land management by farmers along the watershed is made traditionally. Generally, the location of these agricultural activities is on the river border with a moderate to steep slope, so the area has the potential to experience erosion [6, 7]. The land use pattern usually goes on for a long time from generation to generation to form a habit [8]. The slope class category around the Kambaniru watershed is dominated by the high steepness category (slope between 41-60%) with an area of 88,822 hectares (77.92%).

The slope of the land is prone to erosion, especially when the intensity of rainfall is high and vegetation cover is low. The probability of increasing decline is more elevated in line with the pressure on land use for agriculture, gardening, and grazing activities. According to Fitri [7], the sensitivity of soil erosion on forest land, mixed gardens, dry fields, and shrubs along the watershed is generally on a slope. It means the potential for a significant decline can occur if the land use activity gets more prominent and the hill is steeper. Ideally, agricultural business activities on sloping land should be terraced, so they are not prone to erosion [9]. According to Njurumana [2], the criteria for land slope, decay, and runoff threats in the Kambaniru watershed are categorized into three threats, namely low (3.34%), moderate (18.75%), and high (77.92%).

Regarding land management in watersheds, the quality of human resources is closely related to behavior in managing natural resources and the level of dependence on nature. The low rate of human resources causes the knowledge, capacity, and skills to manage natural resources to be quiet, thus having implications for decisions to use processes that can damage the environment. In addition, low human resources cause alternative employment opportunities to become limited and increasingly dependent on nature. As a result of this condition, critical land continues to increase from time to time.

Communities in the upstream and central areas along the Kambaniru watershed are dominated by the livelihoods of farmers (48%), ranchers (5%), fishermen (2%), and traders (4%), with the remaining 41% spread over livelihoods [1]. It shows that the quality of human resources is minimal and depends on nature. Meanwhile, there are more alternative livelihoods in the downstream area of the Kambaniru watershed. It shows that the human resources in the downstream region of the Kambaniru watershed are pretty good, and they are less dependent on nature to support their families.

Based on identifying the problems above, it is crucial to find a solution to prevent erosion of the Kambaniru watershed now and in the future. The effort is made by increasing the capacity of human resources through community empowerment. This study aims to create a causal model of erosion prevention through a human resource development approach to support community food security and develop a technical program for land rehabilitation around the Kambaniru watershed.

2. Method

The data collected consisted of secondary data, including information on environmental issues and damage obtained through data collection and field observations, statistical data, observations, and related research results. The data collected relates to the conditions and problems of damage to land resources in the Kambaniru watershed, covering aspects of land and mineral resources, agriculture, plantations, livestock, and forestry. This data was obtained from relevant agencies in charge of forestry, agriculture, animal husbandry, and plantations.



3. Result and Discussion

3.1. Watershed performance assessment

Based on the Regulation of the Director-General of Land Rehabilitation and Social Forestry Number P.04/V-SET/2009 concerning Guidelines for Monitoring and Evaluation of Watersheds, the results of the evaluation of the performance of the Kambaniru Watershed are moderate. The results of the performance of the Kambaniru watershed in the "moderate" category are the average results of the evaluation of the water system output (water yield), which is in the "fairly good" category, and the value of the condition of the catchment area (DTA), which is in the "almost bad" category. The evaluation value of the water system output is "fairly good," meaning that not all indicators measured are categorized as good. The sedimentation rate indicator shows poor performance and is correlated with the value of "kinda bad" catchment conditions. Poor status indicators also occur in the land use index (IPL), land use capability (KPL), erosion index (IE or CxP), dependence on land (LQ), and land productivity (PL). The performance of the Kambaniru watershed is more emphasized in conditions that have never experienced drought and flooding, including the status of good water quality and low levels of water pollution.

The erosion value is strongly influenced by several indicators such as rainfall, soil erodibility, slope length, slope grade, land cover conditions, and conservation activities, as stated in the USLE formula. The analysis results show that the total erosion rate in the upstream area is 63,769,492.90 ha/ton/year. In contrast, in the middle region, the total erosion rate is 24,034,578.60 ha/ton/year, and in the downstream region, the total erosion rate is 24,034,578.60 ha/ton/year. 4,741,378.55 ha/ton/yr. The erosion index value in the upstream area of the watershed is 700.38%, the middle area is 489.02%, and the downstream area is 276.84%. Based on the classification level of the three river segments, the score is bad, with a score of 5. Land cover and management factors, as well as conservation measures (CP), greatly affect the amount of erosion in the Kambaniru watershed. The CP value in the Upstream area of 0.554, the Middle of 0.633, and the Downstream of 0.547. All three have a score of 5 or are in the Poor classification.

The high sedimentation rate is caused by land cover and poor land suitability. In addition, it is influenced by the habitual patterns of people's behavior so far. Economic conditions and the low level of welfare of the population will affect how land and water are used and managed around the Kambaniru watershed. The results of the performance assessment of the Kambaniru watershed have a score of 3.27 with a moderate classification value (table 1).

3.2. Human resources and pressure on land and forests (upstream, downstream)

Pressure and dependence on land and forests are often associated with community poverty. The same thing happened in the Kambaniru watershed area. Most of the population (> 50%) is below the poverty line. Based on the survey results in the field, the income level of the population per head of household (KK) living in the upper and central areas is on average USD 1–USD 1.5 per day. Downstream is USD 2 per day. Thus, most of the population in the Kambaniru watershed is categorized as poor at around USD 1.0–USD 2.0 per day.

Table 1. Criteria and performance indicators in the Kambaniru watershed

INDICATOR/ PARAMETER	WEIGHT			
	%	%	%	Skor
A. WATER MANAGEMENT	50			0
1. FLOOD AND DROUGHT		30		0
a) River regime coefficient (KRS)			10	3
b) Coef of variance (CV)			5	0
c) Water use index (IPA)			5	0
d) Runoff coefficient (C)			10	3
2. SEDIMENTATION		10		0
- Sedimentation rate			10	5
3. WATER POLLUTION LEVELS		10		0
- Physics (color, TDS, turbidity)			4	1
- Chemical (pH, DHL, NO ₃ , SO ₄ , PO ₄ , K, Ca, Mg)			4	1
- Biology (organic substances, BOD and COD)			2	0
Quantity (A)	50	50		118
Water Management Classification Value				2.36
Category				Fair Good
B. DATA	50			0
1. LAND MANAGEMENT		20		0
a) Land use index (IPL)			4	5
b) Land use capability (KPL)			4	5
c) Erosion index (IE or CxP)			8	5
d) Landslide vulnerability (KTL)			5	3
2. SOCIAL AND ECONOMIC		10		0
a) Individual Concern (KI)			3	1
b) Community participation (PM)			3	1
c) Population pressure (TP)			4	3
3. ECONOMY		10		0
a) Dependence on land (LQ)			4	5
b) Income level (TD)			2	5
c) Land productivity (PL)			2	5
d) Environmental services (JL)			2	3
4. INSTITUTIONS		10		0
a) Empowerment of local institutions (KLL)			2	5
b) Community dependence on the government (KMP)			2	5
c) Coordination, Integration, Synchronization, Synergy (KISS)			4	5
d) Joint business activities (KUB)			2	5
Quantity (B)	50	50		209
Water Management Classification Value				4.18
Category				Kinda Bad
Total (A+B)	100	100		3.27
Watershed Performance Classification Value	100	100	100	3.27
				Moderate

Community poverty is generally caused by minimal natural resources and the low quality of human resources. In addition, climatic conditions with only 3-4 rainy months per year impact the food products produced by the community. Where there is only enough to meet the needs for 3-4 months per year, the low quality of human resources and the absence of other alternative work causes the fulfillment of food needs to only last up to 8-9 months. People only rely on natural conditions by gathering forest products and hunting. It's just that the techniques used tend to damage the environment. For example, they were hunting by burning fields or making their way into the forest by burning without any conservation efforts. The traditional farming system using slash-and-burn is considered by the community around the Kambaniru watershed an easy, cheap, and fast way to grow. Burning is done when hunting animals and clearing fields. Another reason is to produce a fresh, green pasture that is good for animal feed and to allow access to the forest to take tubers, especially gadung (*Dioscorea hispida* Dennst.) tubers. The poor tend to be unable to take adequate adaptation measures due to lacking human resource capacity and accessibility. So, social protection measures, improvement of human resources, and diversification of livelihoods with pro-poor adaptation are necessary [10-12].

3.3. Erosion prevention through HR development

The role of forest and land rehabilitation (RHL) plants can provide direct benefits to the community by increasing food security and employment opportunities and increasing people's income. The RHL plants are types of plants or trees that can be used as a source of food or livelihood that can increase the population's income in the Kambaniru watershed. Planting RHL crops with community involvement through participation has a double impact. First, it aims to improve watershed performance, second, to help the community by providing food, and third, increasing employment in the forestry and plantation product industries [6].

Agricultural food crops such as corn in Nusa Tenggara have a high sustainability index considering that economically, ecologically, and socio-culturally it is very feasible to cultivate. Winarso [13] stated that corn cultivation in Nusa Tenggara is possible to develop because it is supported by regional conditions, technology availability, market conditions, and domestic market demand. On the other hand, using land with steep slopes in the form of hilly and mountainous areas for corn plants, from the aspect of environmental sustainability, is very worrying. Therefore, efforts are needed from the local government to deal with dry land with steep slopes. Socialization and dissemination must be encouraged to provide an understanding of environmental sustainability. For example, they were growing food crops in combination with annual harvests of high economic value. For example, annual plants that are suitable for both lowland (Robusta coffee) and highland (Arabica coffee) can be planted parallel to the contour, then combined with stone piles so that, in the long term, they will form a mound terrace, which functions to hold water and soil from erosion [14]. In addition, given alternatives and types of commodities that can be developed, the community chooses the types of annual plants that interest the community. Tree species are easy to maintain and have high economic value to benefit the community. Besides that, food crops can be cultivated without causing land degradation.

Erosion prevention through capacity building of community human resources in the Kambaniru watershed can be encouraged by planting agricultural land and livestock. A community empowerment program is needed, including outreach activities, training, and mentoring. Through community empowerment programs, people can increase their perceptions, preferences, understanding, and skills to support the success of watershed management. Other positive impacts include job creation for the community. Support by the community in determining the type and suitability of plants will increase the success of planting and plant quality. The usefulness of plant species will increase the Land Use Index (IPL), which will encourage an increase in the performance of the Erosion Index.

Based on figure 1, the causal loop shows that the determination of versatile plant species is not only to increase the erosion index and watershed health but also to support the livelihoods of the people in the Kambaniru watershed. In the figure, it can be identified that the direct causes of the decline in health in the Kambaniru watershed are (1) the problem of poverty as the main factor of pressure on land and forests; and (2) agricultural and livestock systems that are less productive and do not pay attention to conservation issues have an impact on low food security and land damage.

According to Ruijs *et al.* [15]; Moges and Bhat [11]; and Dilshad *et al.* [10] the problem of poverty is caused by the accumulation of the issues in natural conditions that limit food production, education level, and employment. Eventually, there will be an emphasis on land and forests in meeting their food needs. Not only that, the farming and livestock systems practiced by the community, especially in the upstream watershed area, can add pressure to the watershed's health, one of which causes environmental degradation such as erosion. Under these conditions, the land can become less productive, which has implications for low food security [10-12]. Therefore, it is necessary to develop indicators of success in preventing land erosion through the development of human resources to support the food security of the community around the Kambaniru watershed. The forms are: (1) increasing employment opportunities, (2) increasing land productivity, (3) increasing food sources (trees and plants), (4) increasing land cover, (5) increasing land use index, (6) reducing erosion, and (7) improvement of watershed performance. The above activities can run effectively if they are supported by community capacity building, one of which is through community empowerment.




Figure 1. Causal relationship for plant life types and increased erosion index

3.4. Land rehabilitation through program strengthening

Based on the socio-economic conditions and the biophysical environment in the Kambaniru watershed area, community empowerment-based forest development requires a method of utilizing space for plants or trees to be developed to improve community welfare and the quality of forest resources. The commodities that can be produced are multi-purpose tree species. Entities that may be generated can be in the form of wood or non-timber. Some examples of entities are: Group (1) Gum and resin namely *Gaharu* (*Aquilaria* Sp), *Nyatoh* (*Palaquium* spp., *Payena* spp.), and *Kemenyan* (*Styrax benzoin*); (2) Fruits, namely *Durian* (*Durio* spp., especially *D. zibethinus*), Cashew (*Anacardium occidentale*), *Kluwek* or *Kepayang* (*Pangium edule*), Candlenut (*Aleurites moluccana*), Coffee (*Coffea* spp.), Pepper (*Piper nigrum*), Nutmeg (*Myristica fragrans*), Petai (*Parkia speciosa*), Breadfruit; (3) other spices, namely cinnamon (*Cinnamomum* spp.), cloves (*Syzygium aromaticum*), various ginger (*empon-empon*); (4) The woods are *Jeunjing* (*Paraserianthes falcataria*), Teak (*Tectona grandis*), Mahogany (*Swietenia macrophylla*), Gmelina (*Gmelina arborea*); (5) Other plants such as Sandalwood (*Santalum album*), Sago (*Metroxylon sago*).

To develop multi-purpose forest plants and characterize food reserves, it is necessary to apply an agroforestry system. This is in addition to the needs of human life as well as rehabilitating critical land. Furthermore, efforts to rehabilitate critical land require legal instruments, management and utilization systems, and institutional regulations that support the realization of community participation [16, 17]. Therefore, various multi-stakeholder approaches are needed, both technically, regarding the suitability of land types and types of plants. The agroforestry system provides sustainable benefits, and, besides that, it can increase community participation in supporting critical land rehabilitation activities and form solid local institutions [6, 18]. For this reason, the development of agroforestry patterns needs to consider the correct approach pattern, which is following the capabilities of human resources, the environment, and socio-culture, and the selection of the type of commodity to be developed.

In the upstream and middle areas, people's livelihoods depend on subsistence agriculture, horticulture, grazing or pastoralism, and tourism, while in the downstream, agriculture and fishing are the main livelihood options [6, 10]. By considering various factors of land conditions and the community where most of the population lives in agriculture and animal husbandry, the agroforestry that can be developed in the Kambaniru watershed include agrosilviculture, sylvopasture, and agrosilvopasture. Based on the characteristics of the technology, the suitability of each optimization program and land rehabilitation technique is described in table 2 below.

Table 2. Land rehabilitation techniques to prevent erosion around the Kambaniru watershed

No.	Strengthening Program	Land Rehabilitation System	Land Rehabilitation Techniques	Consideration Professional and Empirical
1.	Community Forest (HKM)	• Agrosilviculture	<ul style="list-style-type: none"> • Mixed tree species, namely tree of life (food producer), fruit producer (industry), agarwood tree and luxury wood (sandalwood). • Types of crops, namely food crops (rice, corn, and tubers). • Intercropping system. • Construction of water reservoirs. 	<ul style="list-style-type: none"> • HKM is aimed at improving the welfare of the community through the provision of food, employment and income of the population • Limited agricultural land, so dependent on forest land • Food production is sufficient to support 3-4 months • Types of food crops rice, corn and tubers from the forest • Trees of life (such as breadfruit etc.) to increase food reserves • Types of producing fruits for sale in an effort to increase farmers' income • Embung (Retention basin) is made for water sources for agricultural cultivation during the dry season.
		• Agrosilvopasture	<ul style="list-style-type: none"> • In addition to plant species such as those in the agrosilviculture system, fodder crops such as elephant grass (<i>Pennisetum purpureum</i>) and turi (<i>Sesbania grandiflora</i>) are also planted. 	<ul style="list-style-type: none"> • In addition to the agrosilviculture system, it can also be used for livestock with a cage system.
2.	Village Forest (HD)	• Agrosilviculture	<ul style="list-style-type: none"> • Mixed species of trees of life (food producer), fruit trees producing industrial raw materials (betel nut), agarwood tree, and luxury wood (sandalwood). • Types of crops, namely food crops (rice, corn and tubers). • Intercropping system. • Construction of water reservoirs. 	<ul style="list-style-type: none"> • The Village Government does not have land for village assets or other limited sources of income. • Planting intercropping agricultural crops to meet the food needs of village officials who manage them. • Types of food crops rice, corn and tubers from the forest. • Trees of life (such as breadfruit etc.) to increase food reserves for the villagers. • Embungs are made for water sources for agricultural cultivation during the dry season.
3	Private Forest (HR)	• Agrosilviculture	<ul style="list-style-type: none"> • Tree species are tree species that produce wood for the carpentry industry, industry, as well as essential (aetheric) oils and energy. • Intercropping system with food crops (rice, corn, or tubers). 	<ul style="list-style-type: none"> • HR is aimed at meeting the needs of industrial raw materials, expanding employment and community income. • Demand for wood for industrial raw materials is high. • Land owned for planting community forest is available. • The food crops produced are rice, corn and tubers. • Embungs are used for agricultural cultivation during the dry season
		• Agrosilvopasture	<ul style="list-style-type: none"> • Tree species are tree species that produce wood for the purposes of carpentry, industry, as well as essential oils and energy mixed with leaf-producing plants for animal feed. • Intercropping system with food crops (rice, corn, or tubers) and grasses. 	<ul style="list-style-type: none"> • HR to meet the needs of industrial raw materials, employment and community income. • Demand for wood for industrial raw materials is high. • Land owned for planting community forest is available. • Residents own livestock with a cage system. • Produce food crops, namely rice, corn

No.	Strengthening Program	Land Rehabilitation System	Land Rehabilitation Techniques	Consideration Professional and Empirical
				<ul style="list-style-type: none"> and tubers. • HR provides animal feed in the form of tree leaves and grass (elephant). • Embungs are used for agriculture during the dry season. • Most of the land cover that characterizes the savanna forest ie shrubs, grasslands and trees grows very rarely. • The ecological type of savanna is only overgrown by certain trees, shrubs and grass, so that the types of trees planted are prioritized to be endemic, and those that can increase fertility. • The tree is also used to protect animals and herders
4	Agroforestry and Civil Building Program	<ul style="list-style-type: none"> • Agrosilvopasture 	<ul style="list-style-type: none"> • Tree species are shade trees that are suitable for living in savanna areas • The system of intercropping with food crops (paddy, corn, or tubers) in the valleys (basins). • Planting trees is carried out starting from valley areas and locations that allow them to grow (near water sources). • Construction of water reservoirs. • Strengthening the management of grasslands with a system that does not use fire (burning). 	<ul style="list-style-type: none"> • Most of the people's livelihood is livestock farming with a grazing system. • Tree planting will be successful if it is carried out on soil that has relatively deep top soil, and sufficient soil water content, so that planting is carried out on a limited basis in that location, which is indicated to be in a valley. • Planting of agricultural crops for food crops can be done separately from trees, which allows seasonal crops to be planted. • Water reservoirs are made for water sources for agricultural cultivation and drinking for livestock. • It is necessary to limit the rejuvenation of grass by burning.
		<ul style="list-style-type: none"> • Civil building construction 	<ul style="list-style-type: none"> • Construction of water reservoirs. • Constructed channels to intercept waterways on the slopes. • Build water flow dams in the valleys. 	<ul style="list-style-type: none"> • Rainfall conditions of less than 1500 mm per year prevent water from being stored in the ground. • Prevent water loss by constructing reservoirs such as water reservoirs, interceptor channels, and dams. • The potential for standing water for a long time is used to plant crops around the location of the puddle.
5	Program of Silvopastoral	<ul style="list-style-type: none"> • Silvopasture by expanding shade trees 	<ul style="list-style-type: none"> • Tree species are shade trees that are suitable for living in savanna areas • Not planting agricultural crops because the soil is not suitable for seasonal crops • Planting of trees is carried out starting from valley areas and locations that allow them to grow (near water sources). • Construction of water reservoirs. • Strengthening of grassland management with a system not using fire (burning). 	<ul style="list-style-type: none"> • This system is used in savanna locations where there are very few tree stands or only grassland savanna. • Increase tree planting to curb savanna expansion and desertification.
	Development of Various Forestry Enterprises	<ul style="list-style-type: none"> • Silvopasture by expanding trees that produce non-timber forest products (HHBK /NTPF) 	<ul style="list-style-type: none"> • Types of trees that produce NTFPs that can be used as people's livelihoods, such as calliandra plants and other flowering plants for honey bee business, planting fruit or tubers for food crops, etc. • Planting tree crops is not taken from the wood, but the flowers, fruit and leaves are used for other production. • Planting trees is carried out starting from valley areas and locations that allow them to grow (near water 	<ul style="list-style-type: none"> • Forest honey from East Nusa Tenggara (NTT) is well-known as a non-timber forest product, so it has the potential to be developed in the Kambaniru watershed • The NTFP production process does not make direct use of wood, so the community continues to maintain the presence of the plant or tree. • The required production process is a fast production process. • Commodities produced can increase

No.	Strengthening Program	Land Rehabilitation System	Land Rehabilitation Techniques	Consideration Professional and Empirical
			sources). <ul style="list-style-type: none"> • Construction of water reservoirs. • NTFP business development outside forest areas. 	business and also food security.
6	Environmental Services	<ul style="list-style-type: none"> • Maintaining the existence of Conservation Forests and Nature Preservation for the use of environmental services such as recreation areas and producing drinking water 	<ul style="list-style-type: none"> • Conduct socialization and increase understanding and perception of the importance of conservation areas and nature conservation for the lives of local communities. • Identify environmental services that can be extracted from conservation forests for surrounding communities. • Utilizing environmental services using the principles of sustainability. 	<ul style="list-style-type: none"> • Forest Conservation and Preservation Area is located in the Upper Kambaniru Watershed, so that apart from its main function as nature conservation, it also functions as a water catchment area; • In increasing the existence of Conservation Forests and Nature Preservation Forests, it is necessary to increase the development of Environmental Services for residents around the forests; • Environmental service businesses that may be developed include the development of recreation areas and drinking water businesses.
7	Environmental Greening	<ul style="list-style-type: none"> • Agrosilvopasture 	<ul style="list-style-type: none"> • In private lands around settlements, it is necessary to plant reforestation to support the environment as well as the population's livestock business. • Types of trees planted are trees that can be used as forage for livestock and fruit crops. 	<ul style="list-style-type: none"> • This system is used in residential areas that can support environmental improvement and at the same time support livestock or fruit crops. • Increase the availability of forage trees to support animal feed and increase fruit trees on land around settlements (gardens, fields).

4. Conclusion

The slash-and-burn habit or culture of cultivating land by the community is difficult to eliminate. Therefore, a program to increase the cropping index is needed so that it can improve land quality and productivity. Cultivation technology innovations and supplementary water resources (dams, reservoirs, and others) are the main leverage points in increasing the cropping index, production, and plant productivity. The type of technology needed by the community is not only easy and cheap but also efficient in supporting food crop farming and local government food security programs.

The policy implications of this study are as follows: The limited capacity of community human resources in land conservation can be started by the socialization of planting short-lived trees from the legume family species that can thrive in various types of soil and are flexible with climate and climate change. Corn and tubers are a source of high calories and nutrients needed to improve the quality of human resources in the Kambaniru watershed area. Therefore, developing a food reserve land program in the Kambaniru watershed is necessary.

5. Reference

- [1] BPS-EastSumba 2021 *East Sumba regency in figures 2021* (Waingapu: BPS East Sumba Regency)
- [2] Njurmana G N D 2008 Study of land degradation on kambaniru watershed, East Sumba regency *Info Hutan* **5** 241-54
- [3] Krisnawati H, Varis E, Kallio M and Kanninen M 2011 *Paraserianthes falcataria (L.) Nielsen: ecology, silviculture and productivity*: Cifor)
- [4] Hashim M, Hazim M and Syafinie A 2015 Strategic forest plantation establishment in Malaysia for future product development and utilization *International Journal of Agriculture, Forestry and Plantation* **1** 14-24
- [5] Angima S, Stott D, O'neill M, Ong C and Weesies G 2002 Use of calliandra–Napier grass contour hedges to control erosion in central Kenya *Agriculture, ecosystems & environment* **91** 15-23

- [6] Sukwika T 2019 Participation of community provide hydrological environmental services in the watershed area *Sustainable Environmental and Optimizing Industry Journal* **1** 27-37
- [7] Fitri R 2011 Mapping of erosion hazard levels based on land use and land slope in sub-watersheds Krueng Simpo *Lentera: Jurnal Ilmiah Sains dan Teknologi* **11** 80-6
- [8] Suwarno J 2011 Policy development of sustainable management of upper Ciliwung watershed, district Bogor. (Bogor: IPB University)
- [9] Nyssen J, Poesen J, Haile M, Moeyersons J and Deckers J 2000 Tillage erosion on slopes with soil conservation structures in the Ethiopian highlands *Soil and Tillage Research* **57** 115-27
- [10] Dilshad T, Mallick D, Udas P B, Goodrich C G, Prakash A, Gorti G, Bhadwal S, Anwar M Z, Khandekar N and Hassan S T 2019 Growing social vulnerability in the river basins: Evidence from the Hindu Kush Himalaya (HKH) Region *Environmental Development* **31** 19-33
- [11] Moges D M and Bhat H G 2020 Watershed degradation and management practices in north-western highland Ethiopia *Environmental Monitoring and Assessment* **192** 1-15
- [12] Muoni T, Koomson E, Öborn I, Marohn C, Watson C A, Bergkvist G, Barnes A, Cadisch G and Duncan A 2020 Reducing soil erosion in smallholder farming systems in east Africa through the introduction of different crop types *Experimental Agriculture* **56** 183-95
- [13] Winarso B 2012 Prospects and constraints of corn agribusiness development in West Nusa Tenggara province *Jurnal Penelitian dan Pertanian Terapan* **12** 103-14
- [14] Mulyani A and Mamat H S 2019 Management of dry land with dry climate for maize development in Nusa Tenggara *Jurnal Sumberdaya Lahan* **13** 41-52
- [15] Ruijs A, Dellink R B and Bromley D W 2008 Economics of poverty, environment and natural-resource use: introduction *Frontis* 3-15
- [16] Sukwika T, Febrina L and Mulyawati I 2022 Institutional network of the peat ecosystem restoration plan in Riau Province: Hierarchy and classification approaches. In: *IOP Conference Series: Earth and Environmental Science*: IOP Publishing) p 012019
- [17] Sukwika T and Fransisca L 2021 The policy model for sustainable community forest: A factor analysis *Indonesian Journal of Forestry Research* **8** 135-57
- [18] Sukwika T, Yusuf D N and Suwandhi I 2020 The institutional of local community and stratification of land ownership in surrounding community forests in Bogor *Jurnal Manajemen Hutan Tropika* **26** 59-71