

PAPER NAME

Indonesian indigenous plants as a source of antioxidants to treat gastrointestinal .pdf

WORD COUNT

5273 Words

CHARACTER COUNT

30619 Characters

PAGE COUNT

10 Pages

FILE SIZE

428.1KB

SUBMISSION DATE

Sep 20, 2024 1:16 PM GMT+7

REPORT DATE

Sep 20, 2024 1:17 PM GMT+7

● **1% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 1% Internet database
- 1% Publications database
- Crossref database
- Crossref Posted Content database
- 1% Submitted Works database

● **Excluded from Similarity Report**

- Bibliographic material
- Manually excluded text blocks

Indonesian indigenous plants as a source of antioxidants to treat gastrointestinal disorders

^{1,2,*}Sandrasari, D.A., ^{1,3}Andarwulan, N., ¹Faridah, D.N. and ⁴Dewi, F.N.A.

¹Department of Food Science and Technology, Faculty of Agricultural Technology, IPB University

²Department of Food Tech, Faculty of Food Science and Health, Sahid University, 12780, Jakarta, Indonesia

³Southeast Asian Food and Agricultural Science and Technology (SEAFAST) Center, IPB University

⁴Primate Research Center, IPB University, Bogor, Indonesia

Article history:

Received: 6 December 2020

Received in revised form: 29

January 2021

Accepted: 21 April 2021

Available Online: 3 October

2021

Keywords:

Indonesian indigenous plants,

Antioxidant,

Gastrointestinal disorders

DOI:

[https://doi.org/10.26656/fr.2017.5\(5\).707](https://doi.org/10.26656/fr.2017.5(5).707)

Abstract

Gastrointestinal disorders are the most common illnesses that affect people nowadays. Its prevalence and incidence have increased over the last decades. The majority of the people in Indonesia are still using indigenous plants as medicine to treat these infections or diseases. This study was aimed to determine the potential of Indonesian indigenous plants in treating gastrointestinal disorders. The plants were inventoried through searching articles such as Google Scholar, PubMed, ScienceDirect and Indonesian medicinal plants books. The keyword search term used was “gastrointestinal,” “Indonesian,” “medicinal plants,” “antioxidant activity,” and “phenolic compound.” A total of fifty-one species of indigenous Indonesian plants of 32 families were recorded as being used by the Indonesian population to treat gastrointestinal disorders. Gastrointestinal disorders that heavily utilize indigenous plants are diarrhoea, constipation, gastric ulcer and gastritis. Indonesian indigenous plants containing bioactive compounds such as alkaloids, flavonoids, saponins, tannins, steroids, terpenoids and phenolics that can serve as a source of antioxidants to treat gastrointestinal disorders. There were five Indonesian indigenous plants with the most potential as a source of antioxidants to cope with gastrointestinal disorders: *Zingiber officinale* L, *Annona muricata*, *Phyllanthus niruri*, *Curcuma longa* and *Curcuma xanthorrhiza*.

1. Introduction

Traditional plants medicine has been known for a long time and used by the people of Indonesia. This process has been passed down from generation to generation, especially with the concept of back to nature make traditional medicine more desirable. Indonesian indigenous plants are Indonesian native plants from certain regions/ecosystems, including introduced species from other geographic areas but have evolved with the climate and geographic region of Indonesia (Andarwulan, 2012). Most people think that plants can be used as a traditional medicine because it has a lower side effect than chemical drugs. The original plant is widely used as an alternative medicine for an exogenous antioxidant that has been proven effective for treating a variety of ailments including indigestion (Madikizela *et al.*, 2012). The ability of plants to treat gastrointestinal disorders because of their antioxidant compounds such

as flavonoids, polyphenols, saponins, alkaloids, and terpenoids. Flavonoids and polyphenol compounds are antioxidants, antidiabetic, anticancer, anti-inflammatory and antiseptic, while alkaloids have strong antineoplastic properties also inhibits the growth of cancer cells (Saxena *et al.*, 2012; Xu *et al.*, 2017).

Gastrointestinal disorders are diseases that affect the digestive tract function, such as food and fluid absorption, digestion or excretion. The disorder causes pain, morbidity and can lead to death. Diseases caused by gastrointestinal disorders including 10 major killer diseases in the world. In terms of mortality, causes of death for all age groups diarrheal was 3.5%, ranks 13th out of 22 causes of death, both communicable and non-communicable diseases. WHO (2007) conducted a review of the eight countries in the world, shows that the percentage of occurrences due to gastritis in America by 47%, followed by India at 43% and Indonesia amounted

*Corresponding author.

Email: dinysandra@gmail.com

to 40.85%, gastrointestinal disorders are abdominal or stomach pain, dysentery, gastroenteritis, constipation and vomiting.

The study of Indonesian plants as medicine gastrointestinal disorders have been conducted, but no studies have inventoried the plant as a drug used to treat gastrointestinal disorders. Therefore, this study aimed to identify the Indonesian indigenous plants with traditional benefits in alleviating types of diseases, used parts of the plants and their phytochemical contents. In addition, potential plants as a source of antioxidants in the treatment of gastrointestinal disorders were determined. Plant inventory and determination of potential levels are expected to institutionalize traditional medicine and alternative pharmaceutical preparations.

2. Material and methods

2.1 Inventory of plants

Literature reviews published in journals, reports and books dealing with traditional uses of medicinal plants in Indonesia to treat various gastrointestinal disorders were obtained. Different online databases were used PubMed, Science Direct, Scopus, and Google Scholar with specific search terms such as 'medicinal plants', 'Indonesian plants AND gastrointestinal', phytochemical AND phenolic content AND antioxidant activity from each Indonesian plant were have inventoried. The articles obtained are filtered by title, abstract and full text. Articles that are not relevant to the research topic was issued, while the elected subsequently evaluated using inclusion and exclusion criteria for the study. The inclusion criteria used in this review were articles published in the year 2009 - 2019, Indonesian plants, gastrointestinal, ethanol extract, total phenolic content (TPC) expressed in mg GAE/g extract, total flavonoid content (TFC) expressed in mg QE/g extract and antioxidant activity using DPPH (2,2-diphenyl-1-picrylhydrazyl) method expressed in IC₅₀ with ascorbic acid comparison compound (AEAC) from each of these Indonesian indigenous plants. The term 'Indonesian plants' was used to limit the geographical scope of the search. Selection of ethanol extract and DPPH method because many of these are found in this literature search.

2.2 Phenolic content and antioxidant activity

Total phenolic content and antioxidant activity were used to determine the potential of Indonesian indigenous plants in treating gastrointestinal disorders. The method used is to compare the phenolic content (TPC) with antioxidant activity. Units of data synchronized with the assumption that the ethanol extract of these plants has the same density (ml extract = g of extract). Plants that have the highest potential level value is expressed as the

plant with the most potential as a source of antioxidants to treat gastrointestinal disorders.

2.3 Analysis data

The analysis data consisted of qualitative and quantitative data. Qualitative data includes the results of an inventory of indigenous Indonesian plants such as species names, local names, families, parts used, and types of gastrointestinal disorders that can be treated using these plants. The results of an inventory of the types of gastrointestinal disorders are categorized based on abnormalities or physiological effects of the plant. This categorizing aims to determine the number of plants that can be used for disorders of the gastrointestinal tract. In addition, qualitative data have also used the presence of bioactive compounds such as phenolic, alkaloids, flavonoids, tannins, saponins, terpenoids and steroids.

Quantitative data is used to express the percentage of the number of plants that can be used to treat gastrointestinal disorders. Determine if the potential level of each plant species were analyzed using the XLSTAT program.

3. Result and discussion

3.1 Potential indigenous Indonesian plants

The results of the literature search obtained 97 plants that are known to treat gastrointestinal disorders. Then the articles were selected based on title, abstract and availability of the full-text article. Indonesia's inventory of indigenous plants that can treat gastrointestinal disorders is presented in Table 1. Data in Table 1 shows that 51 species are consisting of 32 plant families that are commonly used to treat gastrointestinal disorders. The most dominant family with respect to a number of species used to treat gastrointestinal disorders is *Zingiberaceae* (ten species) followed by *Apiaceae* (three species), *Solanaceae*, *Fabaceae*, *Piperaceae*, *Annonaceae*, *Arecaceae*, *Punicaceae*, *Acanthaceae*, *basellaceae* and *Myrtaceae* (2 species) whereas other plants 1 species.

To determine the number of plants that can be used to treat gastrointestinal disorders, the types of gastrointestinal disorders are categorized based on abnormalities similar or physiological effects of these plants such as constipation/laxative/constipation, gastritis/gastric ulcer/stomach pain/carminative/flatulence and emetic/nausea/vomiting. Types of gastrointestinal disorders are divided into 13 categories (Table 2). Types of gastrointestinal disorders that can be treated with indigenous Indonesian plants are gastritis (24%), followed by diarrhoea (15.4%), digestive disorders (15.4%), constipation (11.5%), colitis (8.7%),

Table 1. Identified Indonesian indigenous plants in alleviating gastrointestinal disorders

Name	Family	Local name	Parts of Plants Studied	Traditional Uses	Mode of application	References
<i>Acorus calamus</i>	<i>Acoraceae</i>	Jeringau	Leaf	Carminative, choleric, constipation, gastritis, diarrhea	Infusion	Mulyani et al. (2017)
<i>Aloe Vera Linn.</i>	<i>Liliaceae</i>	Lidah buaya	Leaf	Gastritis	Extract, juice	Darwis (2012)
<i>Alpinia galanga</i>	<i>Zingiberaceae</i>	Lengkuas	Rhizome	Diarrhea	Extract, powder	Darwis (2012)
<i>Amomum compactum</i>	<i>Zingiberaceae</i>	Kapulaga	Leaf	Constipation, gastritis, indigestion	Extract, powder	Astra et al. (2019)
<i>Andrographis paniculata Ness.</i>	<i>Acanthaceae</i>	Sambiloto	Leaf	Dysentery, gastric ulcer, sprue	Infusion, extract, powder	Solikhah (2016)
<i>Annona muricata L.</i>	<i>Annonaceae</i>	Sirsak	Fruit	Diarrhea, gastritis	Juice, eat	Abdu (2018)
<i>Annona squamosa L</i>	<i>Annonaceae</i>	Srikaya	Fruit	Indigestion, gastric ulcer	Juice, eat	Rianto et al. (2015)
<i>Anredera cordifolia</i>	<i>Basellaceae</i>	Binahong	Leaf	Dysentery	Infusion, extract	Purwasih et al. (2017)
<i>Areca catechu</i>	<i>Arecaceae</i>	Pinang	Fruit	Indigestion, constipation	Infusion, eat	Darwis (2012)
<i>Averrhoa bilimbi L</i>	<i>Oxalidaceae</i>	Belimbing	Fruit	Indigestion, gastric ulcer	Juice, eat	Darwis (2012)
<i>Caesalpinia sappan L</i>	<i>Caesalpiniaceae</i>	Secang	Stem	Diarrhea	Extract	Widowati (2011)
<i>Camellia sinensis (L.) Kuntze</i>	<i>Theaceae</i>	Teh	Leaf	Diarrhea	Extract	Bedrood et al. (2018)
<i>Carica papaya L.</i>	<i>Caricaceae</i>	Papaya	Fruit	Constipation, gastric ulcer	Juice, eat	Yusro et al. (2019)
<i>Cassia alata L.</i>	<i>Leguminosae</i>	Ketepeng Cina	Leaf	Constipation	Extract, eat	Darwis (2012)
<i>Centella asiatica</i>	<i>Apiaceae</i>	Pegagan	Leaf	Gastritis, carminative	Extract	Darwis (2012)
<i>Cinnamomum cassia Presl</i>	<i>Lauraceae</i>	Kayu manis	Stem	Stomach ache, indigestion	Infusion, extract	Kumar et al. (2019)
<i>Citrus limon</i>	<i>Rutaceae</i>	Lemon	Fruit	Carminative, choleric, constipation, diarrhea, gastric, stomachache, nausea	Juice, eat	Rokaya et al. (2014)
<i>Clitoria ternatea Linn</i>	<i>Fabaceae</i>	Telang	Flower	Stomach ache,	Infusion, extract	Manjula et al. (2013)
<i>Cosmos caudatus</i>	<i>Asteraceae</i>	Kenikir	Leaf	Gastric	Decoction, juice, eat	Cheng et al. (2016)
<i>Curcuma aeruginosa Roxb</i>	<i>Zingiberaceae</i>	Temuireng	Rhizome	Stomach ache	Extract, powder	Suparni (2012)
<i>Curcuma heyneana</i>	<i>Zingiberaceae</i>	Temugiring	Rhizome	Dysentery, constipation	Extract, powder	Subositi and Wahyono (2019)
<i>Curcuma longa</i>	<i>Zingiberaceae</i>	Kunyit	Rhizome	Diarrhea , gastritis, gastric ulcer	Extract, powder	Darwis (2012)
<i>Curcuma xanthorrhiza</i>	<i>Zingiberaceae</i>	Temulawak	Rhizome	Gastritis, diarrhea, dysentery, constipation, gastric ulcer, appetizer	Extract, powder	Darwis (2012)
<i>Curcuma zedoaria</i>	<i>Zingiberaceae</i>	Kunyit putih	Rhizome	Carminative, indigestion, anthelmintic, gastric ulcer, gastritis	Extract, powder	Darwis (2012)
<i>Cyclea barbata</i>	<i>Menispermaceae</i>	Cincau	Leaf	Indigestion, diarrhea, gastritis, ulcer	Extract, gel	Darwis (2012)
<i>Foeniculum vulgare L</i>	<i>Apiaceae</i>	Adas	Fruit	Gastritis	Extract, powder	Darwis (2012)
<i>Guazuma ulmifolia</i>	<i>Malvaceae</i>	Jati Belanda	Leaf	Gastritis	Decoction	Patil and Biradar (2013)

Table 1(Cont.). Identified Indonesian indigenous plants in alleviating gastrointestinal disorders

Name	Family	Local name	Parts of Plants Studied	Traditional Uses	Mode of application	References
<i>Hibiscus sabdariffa</i> L	<i>Mahveaceae</i>	Rosella	Flower	Indigestion, gastric ulcer	Decoction	Suzery et al. (2010)
<i>Jatropha curcas</i> L	<i>Euphorbiaceae</i>	Jarak	Leaf	Gastritis	Oil, poultice	Abdelgadir and Staden (2013)
<i>Kaempferia galanga</i>	<i>Zingiberaceae</i>	Kencur	Rhizome	Stomach ache	Extract, powder	Rao and Kaladhar (2014)
<i>Momordica charantia</i>	<i>Curcubitaceae</i>	Pare	Fruit	Indigestion	Juice, eat	Septimingsih et al. (2017)
<i>Morinda citrifolia</i> L	<i>Rubiaceae</i>	Mengkudu	Fruit	constipation	Extract, juice	Surya et al. (2009)
<i>Moringa oleifera</i>	<i>Moringaceae</i>	Kelor	Leaf	Gastritis, indigestion	Infusion, juice, eat	Sulastrri et al. (2018)
<i>Myristica fragrans</i>	<i>Myristicaceae</i>	Pala	Fruit	Indigestion, diarrhea	Juice, eat	Mulyani et al. (2017)
<i>Nigella sativa</i>	<i>Apiaceae</i>	Jinten hitam	Seed, leaf	Gastric ulcer, indigestion, diarrhea, nausea	Extract, powder	Mulyani et al. (2017)
<i>Ocimum sanctum</i> L.	<i>Ocinaceae</i>	Kemangi	Leaf	Gastritis, gastric ulcer	Infusion, juice, eat	Sopianti dan Sary (2018)
<i>Phyllanthus niruri</i>	<i>Phyllanthaceae</i>	Meniran	Leaf	Gastric ulcer	Extract	Darwis (2012)
<i>Piper crocatum</i>	<i>Piperraceae</i>	Sirih merah	Leaf	Sprue, gastric ulcer	Extract, eat	Prayitno et al. (2016)
<i>Piper nigrum</i> L.	<i>Piperaceae</i>	Merica	Seed,	Indigestion, appetizer, gastric	Extract, powder	Akbar et al. (2014)
<i>Plectranthus amboinicus</i>	<i>Labiatae</i>	Torbangun	Leaf	Gastritis	Extract, juice	Patel et al. (2010)
<i>Pluchea indica</i> (L)	<i>Compositae</i>	Beluntas	Leaf	Gastritis, dysentery, cholic	Infusion, eat	Darwis (2012)
<i>Psidium guajava</i>	<i>Myrtaceae</i>	Jambu biji	Leaf	Diarrhea, dysentery	Juice, eat	Fratwi (2015)
<i>Punica granatum</i> L.	<i>Lythraceae</i>	Delima	Fruit, Seed	Stomach, diarrhea, dysentery, peptic ulcer	Juice, eat	Darwis (2012)
<i>Salaca edulis</i>	<i>Arecaceae</i>	Salak	Fruit	Diarrhea	Juice, eat	Fitrianingsih et al. (2015)
<i>Solanum torvum</i> Swartz	<i>Solanaceae</i>	Takokak	Fruit	Gastric ulcer, stomach ache, gastritis	Extract	Mardiani et al. (2016)
<i>Strobilanthes crispus</i> Bl.	<i>Acanthaceae</i>	Keji Beling	Leaf	Constipation	Decoction	Darwis (2012)
<i>Syzygium polyanthum</i>	<i>Myrtaceae</i>	Salam	Leaf	Gastritis, diarrhea	Extract	Darwis (2012)
<i>Tamarindus indica</i>	<i>Fabaceae</i>	Asam jawa	Fruit	Indigestion, laxative, diarrhea, dysentery, peptic ulcer, constipation	Paste, poultice	Mulyani et al. (2017)
<i>Zingiber cassumanan</i>	<i>Zingiberaceae</i>	Bangle	Rhizome	Constipation, diarrhea	Extract	Mulyani et al. (2017)
<i>Zingiber officinale</i> L	<i>Zingiberaceae</i>	Jahe	Rhizome	Indigestion, nausea, diarrhea, gastritis	Extract, juice, powder	Pairul et al. (2018)
<i>Zingiber zerumbet</i>	<i>Zingiberaceae</i>	Lempuyang	Rhizome	Gastritis, diarrhea, dysentery	Extract, juice, powder	Darwis (2012)

Table 2. Number of plants used to treat gastrointestinal disorders

Types of gastrointestinal disorders	Abbreviation	Number of Plants	Percentage (%)
Gastritis / stomach pain / gastric ulcer	Gas	25	24.0
Diarrhea	Dia	16	15.4
Indigestion	Ind	16	15.4
Constipation/laxative	Con lax	12	11.5
Ulcer	Ulc	9	8.7
Stomachache	Sto	7	6.7
Dysentery	Dis	6	5.8
Appetizer	App	3	2.9
Colic	Col	3	2.9
Emetic/vomit	EmeVom	2	1.9
Carminative	Car	2	1.9
Other disease	Oth dis	2	1.9
Anthelmintic	Ant	1	1.0

abdominal pain (6.7%) and dysentery (5.8%). Some are used to increase appetite, overcome stomach ulcers, nausea, and flatulence.

The results of the literature search also showed that leaves were the most widely used part of the plant with 24 species (44.4%), followed by 13 species of fruit (24.1%), 9 species of rhizomes (16.7%), 3 seeds and flowers each. species (5.6%) and stems of 2 species (3.7%) as can be seen in Figure 1.




Figure 1. Percentage of plant parts used to treat gastrointestinal disorders

Indigenous Indonesian plants have many bioactive compounds such as phenolic, flavonoids, terpenoids, alkaloids, tannin, terpenoids, saponins and steroids so it can be used to treat several common diseases including gastrointestinal disease. Data bioactive compounds in each plant were extracted using ethanol can be seen in Table 3.

3.2 Phenolic content, flavonoid content and antioxidant activity of identified indigenous Indonesian plants

The results of the literature search on phenolic content (TPC), flavonoid content (TFC) and antioxidant activity are presented in Table 4. Data in Table 4 shows that *Zingiber officinale* is a plant that has the highest potential level as a source of antioxidant to treat gastrointestinal disorders followed by *Annona muricata*, *Phyllanthus niruri*, *Curcuma longa* and *Curcuma xanthorrhiza*.

4. Conclusion

This study clearly shows the potential of Indonesian indigenous plants as antioxidant agents to treat gastrointestinal disorders. Pharmacological studies reveal that a diverse group of naturally occurring chemical substances derived from the plants show promising antioxidant activity such as *Zingiber officinale*, *Annona muricata*, *Curcuma longa* and others. Therefore, further research should be carried out on the bioactive compounds present in the particular plants which have the potential as an antioxidant agent, especially to treat gastrointestinal disorders.

References

- Abdu, S.B. (2018). *Annona muricata* (Graviola): A review of its traditional uses and pharmacological activities. *International Journal of Life Sciences Research*, 6(1), 50-57.
- Abdulkadir, A.R., Mat, N., Hasan M.M. and Jahan, M.S. (2016). *In vitro* antioxidant activity of the ethanolic extract from fruit, stem, and leaf *Solanum torvum*. *Science Asia*, 42(3), 184-189. <https://doi.org/10.2306/scienceasia1513-1874.2016.42.184>
- Agustina, S., Ruslan and Wiraningtyas, A. (2016). Skrining fitokimia tanaman obat di kabupaten Bima. *Cakra Kimia Indonesian. E-journal of Applied Chemistry*, 4(1), 71-76.
- Alaga, T.O., Edema, M.O., Atayese, A.O. and Bankole, M.O. (2014). Phytochemical and in vitro antibacterial properties of *Hibiscus sabdariffa* L. (Roselle) juice. *Journal of Medicinal Plant Research*, 8(6), 339-344. <https://doi.org/10.5897/JMPR12.1139>
- Andarwulan, N., Dewi, K., Riza, A.A., Hardianzah, R., Anna, V.R. and Bradley, W.B. (2012). Polyphenols, carotenoids, and ascorbic acid in underutilized medicinal vegetables. *Journal of Functional Foods*, 4(1), 339-347. <https://doi.org/10.1016/>

Table 3. Phytochemical content of Indonesian indigenous plants

Types of plants	Alkaloids	Flavonoids	Tannins	Saponins	Terpenoids	Steroid	Phenolic	References
<i>Aloe Vera</i> Linn	+	+	+	+	+	+	+	Jha et al. (2018)
<i>Alpinia galanga</i>	+	+	-	+	+	+	++	Agustina et al. (2016)
<i>Amomum compactum</i>	+	+	-	+	+	+	+	Silalahi (2017)
<i>Andrographis paniculata</i> Ness	+	+	+	+	NA	NA	+	Rajalakshmi and Cathrine (2016)
<i>Annona muricata</i>	+	+	+	+	+	+	+	Ojezele et al. (2016)
<i>Annona squamosa</i>	+	+	NA	+	+	+	+	Mulyani et al. (2013)
<i>Averrhoa bilimbi</i> L	++	++	+++	+++	-	+++	+	Hasim et al. (2019)
<i>Caesalpinia sappan</i>	+	+	-	+	+	-	++	Widowati (2011)
<i>Camellia sinensis (L.) Kuntze</i>	+	+	+	+	+	+	+	Martono and Setiyono (2014)
<i>Carica papaya</i>	+	+	+	-	NA	+	+	Zahra et al. (2017)
<i>Cinnamomum cassia</i> Presl	+	+	+	+	+	+	+	Kumar et al. (2019)
<i>Citrus limon</i>	+	+	+	-	NA	+	+	Mayasari and Laoli (2018)
<i>Clitoria ternatea</i> Linn	+	+	+	-	+	-	-	Manjula et al. (2013)
<i>Cosmos caudatus</i>	+++	+	+	+++	+	+	+	Liliwirianis et al. (2011)
<i>Curcuma aeruginosa</i> Roxb	-	+	-	++	+	-	+	Nurcholis (2017)
<i>Curcuma heyneana</i>	++	++	++	+	+	-	++	Ghasemzadeh et al. (2016)
<i>Curcuma longa</i>	+	+	+	+	+	+	+	Agustina et al. (2016)
<i>Curcuma xanthorrhiza</i>	+	+	-	-	-	-	+	Agustina et al. (2016)
<i>Curcuma zedoaria</i>	+	+	-	+	+	+	+	Sumathi et al. (2013)
<i>Cyclea barbata</i>	-	+	+	+	+	+	+	Mentari et al. (2019)
<i>Foeniculum vulgare</i> L	+	+	+	+	-	+	+	Effendi et al. (2015)
<i>Hibiscus sabdariffa</i> L	++	++	++	+++	+	++	++	Alaga et al. (2014)
<i>Kaempferia rhizoma</i>	+	+	+	-	-	NA	+	Hayati et al. (2015)
<i>Momordica Charantia</i>	+	+	+	-	NA	+	+	Septingsih et al. (2017)
<i>Morinda citrifolia</i>	+	++	+	+	++	++	+	Kumar et al. (2014)
<i>Moringa oleifera</i>	+	+	+	+	NA	NA	+	Sulastri et al. (2018)
<i>Myristica fragrans</i>	+	+	+	+	NA	+	+	Gupta et al. (2014)
<i>Nigella sativa</i>	+++	+++	++	-	NA	++	+++	Zarai et al. (2013)
<i>Ocimum sanctum</i> L.	+	+	+	+	+	NA	+	Sopianti and Sary (2018)
<i>Phyllanthus niruri</i>	+	+	+	+	+	NA	++	Ramandeep et al. (2017)
<i>Piper crocatum</i> Ruiz & Pav	++	++	++	++	NA	-	++	Prayitno et al. (2016)
<i>Piper nigrum</i> Linn	+++	++	++	-	+	++	+	Nahak and Sahu (2011)
<i>Psidium guajava</i>	+	+	+	-	-	+	+	Agustina et al. (2016)
<i>Punica granatum</i> L	-	+	+	+	-	+	+	Agustina et al. (2016)
<i>Salaca edulis</i>	+	+	+	NA	+	-	+	Fitrianiingsih et al. (2015)
<i>Solanum torvum</i>	+	+	+	+	NA	NA	NA	Rahman et al. (2013)

Table 3. Phytochemical content of Indonesian indigenous plants

Types of plants	Alkaloids	Flavonoids	Tannins	Saponins	Terpenoids	Steroid	Phenolic	References
<i>Syzygium polyanthum</i>	+	+	+	-	-	+	+	Agustina et al. (2016) ²
<i>Tamarindus indica</i>	-	+	+	+	-	NA	+	Buanasari et al. (2018)
<i>Zingiber cassumunar</i>	+	+	+	+	+	-	+	Buldani et al. (2017)
<i>Zingiber officinale</i>	+	+	-	+	+	+	+	Agustina et al. (2016)

NA, Not available.

Table 4. Total phenolic content (TPC), antioxidant activity and potential levels of indigenous Indonesian plants

Plant species	Total Phenolic Content (TPC)		Total Flavonoid Content (TFC)		Antioxidant Activity (AA)		Potential level (TPC/AA)	References
	(mg GAE/g extract)	(mg QE/g extract)	(mg QE/g extract)	(IC ₅₀) µg/mL	(AA)	(TPC/AA)		
<i>Aloe vera</i> Linn	7.99	9.17	45.6	175.22	+	175.22	Vidic et al. (2014)	
<i>Alpinia galanga</i>	37.84	NA	93.27	405.70	+	405.70	Da'i et al. (2013)	
<i>Annona muricata</i>	195.05	64.25	21.03	9274.85	+	9274.85	Yen et al. (2019)	
<i>Avorhoa bilimbi</i> L	39.03	97.28	16.99	2297.23	+	2297.23	Hasim et al. (2019)	
<i>Camellia sinensis</i>	31.6	23.2	7.30	4328.77	+	4328.77	Bizuayebu et al. (2016)	
<i>Cosmos caudatus</i>	149.33	214.00	166.25	898.23	+	898.23	Noriham et al. (2015)	
<i>Curcuma aeruginosa</i> Roxb	51.49	33.36	406.52	126.66	+	126.66	Kumar et al. (2013)	
<i>Curcuma longa</i>	6.15	4.28	1.08	5694.81	+	5694.81	Wandita and Musfiroh (2018)	
<i>Curcuma xanthorrhiza</i>	205.86	NA	36.96	5569.81	+	5569.81	Amelinda et al. (2018)	
<i>Curcuma zedoaria</i>	84.15	NA	210.12	400.49	+	400.49	Marliani et al. (2017)	
<i>Foeniculum vulgare</i>	5.8	16.8	2.409	2407.64	+	2407.64	Martati and Akmalina (2018)	
<i>Hibiscus sabdariffa</i> L	48.42	2.075	281.78	171.85	+	171.85	Jung et al. (2013)	
<i>Kaempferia galanga</i>	23.85	NA	13.07	1824.79	+	1824.79	Latifah (2015)	
<i>Morinda citrifolia</i>	88.03	1.56	398.99	22.06	+	22.06	Kumar et al. (2014)	
<i>Moringa oleifera</i>	30.0	96.0	134.5	223.05	+	223.05	Sulastri et al. (2018)	
<i>Myristica fragrans</i>	37.26	51.97	770	48.39	+	48.39	Liu et al. (2008)	
<i>Nigella sativa</i>	31.15	110.5	53.90	577.83	+	577.83	Zarai et al. (2013)	
<i>Ocimum sanctum</i> L.	148.9	18.27	52.68	2826.50	+	2826.50	Erviana et al. (2011)	
<i>Phyllanthus niruri</i>	61.36	54.72	10.53	5827.16	+	5827.16	Ramandeep et al. (2015)	
<i>Piper crocatum</i>	182.68	197.95	82.71	2208.68	+	2208.68	Prayitno et al. (2016)	
<i>Piper nigrum</i> Linn	174.92	62.73	448.20	390.27	+	390.27	Nahak and Sahu (2011)	
<i>Plectranthus amboinicus</i>	11.6	1.612	247.94	46.79	+	46.79	Patel et al. (2010)	
<i>Solanum torvum</i>	16.15	1.41	124	130.24	+	130.24	Abdulkadir et al. (2016)	
<i>Syzygium polyanthum</i>	69.76	270	35.057	1989.73	+	1989.73	Verawati et al. (2017)	
<i>Tamarindus indica</i>	71.68	NA	98.30	729.20	+	729.20	Yeasmen and Islam (2015)	
<i>Zingiber cassumunar</i>	27.1	NA	142.7	189.91	+	189.91	Marliani et al. (2015)	
<i>Zingiber officinale</i> L	400.2	268.2	39.6	10106.06	+	10106.06	Yesiloglu et al. (2012)	

Value of the potential level is calculated assuming the density of the extract for each plant is the same.

- j.jff.2012.01.003
- Amelinda, E., Widarta, W.R. and Damayanti, L.P.T. (2018). Pengaruh waktu maserasi terhadap aktivitas antioksidan ekstrak rimpang temulawak. *Jurnal Ilmu dan Teknologi Pangan*, 7(4), 165-174. <https://doi.org/10.24843/itepa.2018.v07.i04.p03> [In Bahasa Indonesia].
- Asra, R., Azni, N.R., Rudi and Nessa. (2019). Uji aktivitas antioksidan ekstrak etanol fraksi heksan, etil asetat dan air daun kapulaga. *Journal of Pharmaceutical and Sciences*, 2(1), 30-37. <https://doi.org/10.36490/journal-jps.com.v2i1.17> [In Bahasa Indonesia].
- Bedrood, Z., Rameshrad, M. and Hosseinzadeh, H. (2018). Toxicological effect of *Camellia sinensis* (green tea): A review. *Phytotherapy Research*, 32(7), 1163 – 1180. <https://doi.org/10.1002/ptr.6063>
- Bizuayebu, D., Atlabachew, M. and Ali, M. (2016). Determination of some selected secondary metabolites and their in-vitro antioxidant activity in commercially available Ethiopian tea (*Camellia sinensis*). *Springer Plus*, 5, 412. <https://doi.org/10.1186/s40064-016-2056-1>
- Cheng, S.H., Barakatun-Nisak, M.Y., Anthony, J. and Ismail, A. (2016). Potential medicinal benefits of *Cosmos caudatus* (Ulam Raja): A scoping review. *Journal of Research in Medicinal Sciences*, 20(10), 1000-1006. <https://doi.org/10.4103/1735-1995.172796>
- Da'i, M., Wahyuni, A.S., Trisharyanti, I., Azizah, T., Suhendi, A. and Saifudin, A. (2015). Antioxidant activity of *Phyllanthus niruri* L. Herb: *in vitro* and *in vivo* models and isolation of active compound. *National Journal of Physiology, Pharmacy and Pharmacology*, 6(3), 32-37. <https://doi.org/10.5455/njppp.2015.5.0510201575>
- Darwis, W. (2012). Tanaman obat yang terdapat di kota Bengkulu yang berpotensi sebagai obat penyakit dan gangguan pada sistem pencernaan manusia. *Konservasi Hayati*, 8(1), 1-15. [In Bahasa Indonesia].
- Fratiwi, Y. (2015). The Potential of guava leaf (*Psidium guajava* L.) for diarrhea. *Journal of Majority*, 4(1), 113-118.
- Ghasemzadeh, A., Jaafar, H.Z.E. and Rahmat, A. (2016). Variation of the phytochemical constituents and antioxidant activities of *Zingiber officinale* var *rubrum* theilade associated with different drying methods and polyphenol oxidase activity. *Molecules*, 2(1), 780-788. <https://doi.org/10.3390/molecules21060780>
- Gupta, A.D., Vipin, K.B., Vikash, B. and Nishi, M. (2013). Chemistry, antioxidant and antimicrobial potential of nutmeg (*Myristica fragrans* Houtt). *Journal of Genetic Engineering and Biotechnology*, 11(1), 25-31. <https://doi.org/10.1016/j.jgeb.2012.12.001>
- Jha, Prakash, D. and Bisht, D. (2018). A phytochemical screening of the ethanolic extract of Aloe vera gel. *International Journal of Science and Research*, 8 (10), 1543-1545.
- Kumar, S. and Pandey, A. (2013). Chemistry and Biological Activities of Flavonoids: An Overview. *The Scientific World Journal*, 1-16. <https://doi.org/10.1155/2013/162750>
- Kumar, K., Suresh, M., Kumar, S.A. and Kalaiselvi, P. (2014). Bioactive compounds, radical scavenging, antioxidant properties and FTIR spectroscopy study of *Moringa citrifolia* fruit extracts. *International Journal of Current Microbiology and Applied Science*, 3(2), 28-42.
- Kumar, S., Kumar, R. and Mishra, S. (2019). Pharmacological properties and their medicinal uses of *Cinnamomum*: A review. *Journal of Pharmacy and Pharmacology*, 71(12), 1735 – 1761. <https://doi.org/10.1111/jph.13173>
- Liliwirianis, N., Musa, N.L.N.W., Zain, W.Z.W.M., Kassim, J. and Karim, S.A. (2011). Preliminary studies on phytochemical screening of ulam and fruit from Malaysia. *E-Journal of Chemistry*, 8(1), 285-288. <https://doi.org/10.1155/2011/464595>
- Madikizela, B., Ndhlala, A.R., Finnie, J.F. and van Staden, J. (2012). Ethnopharmacological study of plants from Pondoland used against diarrhea. *Journal of Ethnopharmacology*, 141(1), 61-71. <https://doi.org/10.1016/j.jep.2012.01.053>
- Manjula, P., Mogan, C.H., Sreekanth, D., Keerthi, B. and Prathibha, D. (2013). Phytochemical analysis of *Clitoria ternatea* Linn, a valuable medicinal plant. *Journal of Indian Botanical Society*, 92(3 and 4), 173 – 178.
- Martono, B. and Setiyono, R.T. (2014). Phytochemical screening of six tea genotypes. *Journal of Industrial and Beverage Crops*, 1(2), 63-68. <https://doi.org/10.21082/jtidp.v1n2.2014.p63-68>
- Mayasari, U. and Laoli, M. T. (2018). Karakterisasi simplisia dan skrining fitokimia daun jeruk lemon (*Citrus limon* (L.) Burm.f.). *Klorofil*, 2(1), 7-13. [In Bahasa Indonesia].
- Mulyani, M., Arifin, B. and Nurdin, H. (2013). Uji antioksidan dan isolasi senyawa metabolit sekunder dari daun srikaya (*Annona squamosa* L). *Jurnal Kimia Unand*, 2(1), 6-12. [In Bahasa Indonesia].
- Mulyani, H., Harti, W.S. and Indria, E.V. (2017). Pengobatan tradisional jawa dalam manuskrip serat

- primbon jampi Jawi. *Jurnal Literia*, 16(1), 135-151. <https://doi.org/10.21831/ltr.v16i1.14257> [In Bahasa Indonesia].
- Nahak, G. and Sahu, R.K. (2011). Phytochemical evaluation and antioxidant activity of *Piper cubeba* and *Piper nigrum*. *Journal of Applied Pharmaceutical Science*, 1(8), 153-157.
- Nurcholis, W. and Bintang, M. (2017). Perbandingan aktivitas antioksidan dan kandungan fenolik temulawak dan temu ireng. *Jurnal Jamu Indonesia*, 2 (1), 25-29. <https://doi.org/10.29244/jji.v2i1.27> [In Bahasa Indonesia].
- Ojezele, O.J., Ojezele, M.O. and Adeosun, A.M. (2016). Comparative phytochemistry and antioxidant activities of water and ethanol extract of *Annona muricata* Linn leaf, seed and fruit. *Advances in Biological Research*, 10(4), 230-235.
- Patel, R.D., Naveen, K.M., Manjul, P.S., Anita, S., Naheed, W.S., Gulzar, A. and Sudarshan, K.S. (2010). Antioxidant potential of leaves of *Plectranthus amboinicus* (lour) spreng. *Der Pharmacia Lettre*, 2(4), 240-245. [https://doi.org/10.1016/S0975-3575\(10\)80057-4](https://doi.org/10.1016/S0975-3575(10)80057-4)
- Patil, A.B. and Jadhav, A.S. (2013). Flavonoid and antioxidants: A review. *International Journal of Pharmaceutical and Biological Sciences Research and Development*, 1(2), 7-20.
- Prayitno, S.A., Kusnadi, J. and Murtini, E.S. (2016). Antioxidant activity of red betel leaves extract (piper crocatum ruiz and pav.) By difference concentration of solvents. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 7(5), 1836 – 1843.
- Rajalakshmi, V. and Catherine, L. (2016). Phytochemical screening and antimicrobial activity of ethanolic extract of *Andrographis paniculate*. *Journal of Pharmacognosy and Phytochemistry*, 5 (2), 175-177.
- Ramandeep, K., Nahid, A., Neelabh, C. and Navneet, K. (2017). Phytochemical screening of *Phyllanthus niruri* collected from Kerala region and its antioxidant and antimicrobial potential. *Journal of Pharmaceutical Science and Research*, 9(5), 1312-1316.
- Rianto, L., Astuti, I., Handayani. and Septiani, A. (2015). Uji aktivitas ekstrak etanol 96% biji srikaya (*Annona squamosa* l) sebagai antidiare dengan metode difusi cakram. *Jurnal Ilmiah Manuntung*, 1(2), 181-186. <https://doi.org/10.51352/jim.v1i2.33> [In Bahasa Indonesia].
- Rokaya, B.M., Yadav, U., Poudel, R.C., Timsina, B., Münzbergová, Z., Asselin, H., Tiwari, A., Shrestha, S. and Sigdel, S.R. (2014). Traditional uses of medicinal plants in gastrointestinal disorders in Nepal. Review. *Journal of Ethnopharmacology*, 158 (Part A), 221-229. <https://doi.org/10.1016/j.jep.2014.10.014>
- Saxena, N., Shrivastara, P.N. and Saxena, C. (2012). Preliminary physicochemical study of stem bark of *Alstonia scholaris* (L) medicinal plant. *International Journal of Pharmaceutical Sciences and Research*, 3 (4), 1071-1075.
- Silalahi, M. (2017). Bioaktivitas *Amomum compactum Soland ex Maton* dan perspektif konservasinya. *Journal Pro-Life*, 4(2), 320-328. [In Bahasa Indonesia].
- Solikhah, E.N. (2016). Indonesian medicinal plants as sources of secondary metabolites for pharmaceutical industry. *Journal of Medicinal Science*, 48(4), 226-239. <https://doi.org/10.19106/JMedSci004804201606>
- Sopianti. and Sary. (2018). Skrining fitokimia dan profil KLT metabolit sekunder dari daun ruku-ruku (*Ocimum tenuiflorum* l.) dan daun kemangi (*Ocimum sanctum* l). *Jurnal Farmasi dan Kesehatan*, 8(1), 44-52. <https://doi.org/10.36434/scientia.v8i1.118> [In Bahasa Indonesia].
- Sulastri., Zubair, M.S., Anas, N.I., Abidin, S., Hardani, R., Yulianti, R. and Aliyah. (2018). Total phenolic, total flavonoid, quercetin content and antioxidant activity of standardized extract of *Moringa oleifera* leaf from regions with different elevation. *Pharmacognosy Journal*, 10(6), 104-108. <https://doi.org/10.5530/pj.2018.6s.20>
- Sumathi, S., Iswariya, G.K., Sivaprabha, B., Dharani, P., Radha. and Padma, P.R. (2013). Comparative study of radical scavenging activity and phytochemical analysis of fresh and dry rhizomes of *Curcuma zedoria*. *International Journal of Pharmaceutical Sciences and Research*, 4(3), 1069-1073.
- Suzery, M., Lestari, S. and Cahyono B. (2010). Penentuan total antosianin dari kelopak bunga rosella (*Hibiscus sabdariffa* L.) dengan metode maserasi dan sokhletasi. *Jurnal Sains dan Matematika*, 18(1), 1-6. [In Bahasa Indonesia].
- Vidic, D., Tarić, E., Alagić, J. and Maksimović, M. (2014). Determination of total phenolic content and antioxidant activity of ethanol extracts from *Aloe* spp. *Bulletin of the Chemists and Technologists of Bosnia and Herzegovina*, 42, 5-10.
- WHO (World Health Organization). (2007). Traditional Medicine Fact Sheet 134. Geneva: World Health Organization.
- Widowati, W. (2011). Uji fitokimia dan potensi

- antioksidan ekstrak etanol kayu secang (*Caesalpinia sappan* L.). *Jurnal Kedokteran Maranatha*, 11(1), 23-31. [In Bahasa Indonesia].
- Xu, D.P., Li, Y., Meng, X., Zhou, T., Zheng, J., Zhang, J. and Li, H-B.. (2017). Natural antioxidant in foods and medicinal plants: Extraction, assessment and resources. *International Journal of Molecular Science*, 18(1), 96. <https://doi.org/10.3390/ijms18010096>
- Yeasmen, N. and Islam, N. (2015). Ethanol as a solvent and hot extraction technique preserved the antioxidant properties of tamarind (*Tamarindus indica*) seed. *Journal of Advanced Veterinary and Animal Research*, 2(3), 332-337. <https://doi.org/10.5455/javar.2015.b103>
- Yesiloglu, Y., Aydin, H. and Ismail, K. (2012). In vitro antioxidant activity of various extracts of ginger (*Zingiber officinale* L.) seed. *Asian Journal of Chemistry*, 25(7), 3573-3578. <https://doi.org/10.14233/ajchem.2013.13657>
- Yusro, F., Mariani, Y. and Wardenaar, E. (2019). Utilization of medicinal plants to treat digestive system disorders by Dayak Iban Tribe: case study in Kapuas Hulu Regency, West Kalimantan. *Jurnal Borneo Akcaya*, 5(1), 58-72. <https://doi.org/10.51266/borneoakcaya.v5i1.120>
- Zarai, Z., Boujelbene, E., Salem, N.B., Gargouri, Y. and Sayari, A. (2013). Antioxidant and antimicrobial activities of various solvent extract, piperine and piperic acid from *Piper nigrum*. *Food Science and Technology*, 50(2), 634-641. <https://doi.org/10.1016/j.lwt.2012.07.036>
- Zahra, Shaheed, S. and Malik, A. (2017). *In vitro* phytochemical screening and antioxidant activity of *Carica papaya* plant parts collected from Lahore, Pakistan. *Journal of Natural Product and Plant Resource*, 7(4), 23-28.

● 1% Overall Similarity

Top sources found in the following databases:

- 1% Internet database
- Crossref database
- 1% Submitted Works database
- 1% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Universitas Sultan Ageng Tirtayasa on 2019-09-25 Submitted works	<1%
2	myfoodresearch.com Internet	<1%

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded text blocks

EXCLUDED TEXT BLOCKS

Food Research 5 (5) : 195 - 204 (October 2021)Journal homepage: <http://www.myfoodresearch.com>

of Food Tech, Faculty of Food Science and Health, Sahid University, 12780, Jakarta...
www.myfoodresearch.com

eISSN: 2550-2166 / © 2021 The Authors. Published by Rynnye Lyan ResourcesEmail Universitas Nusa Cendana on 2023-04-11

et al. / Food Research 5 (5) (2021)
Universitas Nusa Cendana on 2023-04-11

to 40.85%, gastrointestinal disorders are abdominal orstomach pain, dysentery, ga...
www.myfoodresearch.com

eISSN: 2550-2166© 2021 The Authors. Published by Rynnye Lyan Resources
Padjadjaran University on 2023-05-20

Sandrasari et al. / Food Research 5 (5) (2021) 195 - 204
www.myfoodresearch.com

Table 1. Identified Indonesian indigenous plants in alleviating gastrointestinal diso...
www.myfoodresearch.com

Extract, powderDarwis (2012)Curcuma xanthorrhizaZingiberaceaeTemulawakRhi...
www.myfoodresearch.com

gastric ulcer
www.myfoodresearch.com

Extract, powderDarwis (2012)Cyclea barbataMenispermaceaeCincauLeafIndigesti...

www.myfoodresearch.com

Carminative, indigestion,anthelmintic, gastric ulcer, gastritis

www.myfoodresearch.com

Gastritis, diarrhea, dysentery,constipation

www.myfoodresearch.com

Curcuma longaZingiberaceaeKunyitRhizomeDiarrhea , gastritis, gastric ulcer

www.myfoodresearch.com

eISSN: 2550-2166© 2021 The Authors. Published by Rynnye Lyan Resources

Padjadjaran University on 2023-05-20

Curcuma aeruginosa RoxbZingiberaceaeTemuirengRhizomeStomach ache

www.myfoodresearch.com

Clitoria ternatea LinnFabaceaeTelangFlowerStomach ache

www.myfoodresearch.com

Guazuma ulmifolia

www.myfoodresearch.com

et al. / Food Research 5 (5) (2021)

Universitas Nusa Cendana on 2023-04-11

Table 1

www.myfoodresearch.com

Pluchea indica (L

Eni Nuraeni, Maulana Alkandahri, Sri Mulyanthy Tanuwidjaja, Khuzaimah Nurul Fadhilah et al. "Ethnopharma..."

Table 2. Number of plants used to treat gastrointestinal disordersTypes of gastroi...

www.myfoodresearch.com

4. ConclusionThis study clearly shows the potential of Indonesian indigenous plant...

www.myfoodresearch.com

Figure 1. Percentage of plant parts used to treat gastrointestinal disordersIndigeno...

www.myfoodresearch.com

Effendi et al. (2015)Alaga et al. (2014)Hayati et al. (2015)Septiningsih et al. (2017)...

www.myfoodresearch.com

Prayitno et al. (2016)Nahak and Sahu (2011)Agustina et al. (2016)Agustina et al. (...)

www.myfoodresearch.com

Table 3. Phytochemical content of Indonesian indigenous plants

www.myfoodresearch.com

Curcuma aeruginosa

Noura S. Dosoky, William N. Setzer. "Chemical Composition and Biological Activities of Essential Oils of Cur...

Solanum torvum++++NANA

www.myfoodresearch.com

NA, Not available.Table 4. Total phenolic content (TPC), antioxidant activity and p...

www.myfoodresearch.com

Table 3. Phytochemical content of Indonesian indigenous plantsTypes of plantsAl...

www.myfoodresearch.com

Sulastri et al. (2018)Gupta et al. (2014)Zarai et al. (2013

www.myfoodresearch.com

Sopianti and Sary (2018)Ramandeep et al. (2017

www.myfoodresearch.com